

Patient Enrollment Form | Rare Blood Disorders



NovoCare[®]
Savings | Coverage | Support

Phone: 1-844-668-6732
Fax: 1-866-488-6576

Monday - Friday
8:00 AM to 8:00 PM ET

* Indicates a required field New start Reauthorization Restarting treatment Transitioning from: _____

Page 1 of 3

SERVICES REQUESTED	Select a product: <input type="checkbox"/> NovoSeven [®] RT <input type="checkbox"/> Novoeight [®] <input type="checkbox"/> Tretten [®] <input type="checkbox"/> Rebiny [®] <input type="checkbox"/> Esperoct [®]
	<input type="checkbox"/> Trial Program ^a <input type="checkbox"/> Patient has commercial prescription coverage, such as an HMO or PPO <input type="checkbox"/> Patient has been newly prescribed a Novo Nordisk factor product for an indicated condition <input type="checkbox"/> Patient is Novo Nordisk factor product-naïve <input type="checkbox"/> JumpStart [™] request. Shipping schedule to be confirmed with patient by NovoCare [®] . <input type="checkbox"/> NovoCare [®] Savings Offer (if eligible) For complete copay terms and conditions, visit NovoFactorSavings.com
<small>^a Patients who have been prescribed one of the above products for an FDA-approved indication and who have commercial insurance may be eligible to receive a limited supply of free product from Trial and/or JumpStart[™]. Patient is not eligible if he/she participates in or seeks reimbursement or submits a claim for reimbursement to any federal or state health care program with prescription drug coverage, such as Medicaid, Medicare, Medigap, VA, DOD, TRICARE, or any similar federal or state health care program. Trial and JumpStart[™] products are provided at no cost to the patient or the HCP, is not contingent on any product purchase, and the patient and HCP must not: (1) bill any third party for the free product, or (2) resell the free product. No purchase necessary.</small>	

PATIENT/INSURANCE INFORMATION	Patient name: *	DOB (MM/DD/YYYY):*			
	Gender [†] :* <input type="checkbox"/> Male <input type="checkbox"/> Female	Preferred language: <input type="checkbox"/> English <input type="checkbox"/> Spanish <input type="checkbox"/> Other:			
	Home address (No P.O. box):	City:	State:	Zip:*	
	Shipping address (If different from Home Address):	City:	State:	Zip:*	
	Email:	Primary phone:	Ship drug to: <input type="checkbox"/> Patient's home <input type="checkbox"/> Prescribing HCP		
	Primary guardian/caregiver (required if patient under 18 years old):*	Relationship to patient:			
	Primary pharmacy insurance: (Please attach a copy of the insurance card if available)				Phone:
	Rx # ID:	Rx Group #:	Rx PCN #:	Rx BIN #:	
	Secondary pharmacy insurance: (Please attach a copy of the insurance card if available)				Phone:
	Rx # ID:	Rx Group #:	Rx PCN #:	Rx BIN #:	
<small>[†] Novo Nordisk and its partners recognize that patients may not identify as male or female. However, many insurance companies still require that one of these two fields be used for each of their members. Please indicate the gender on file with the patient's insurance company.</small>					

DIAGNOSIS	What is the primary diagnosis for which you are prescribing a Novo Nordisk factor product? (required)*
	<input type="checkbox"/> D66 - Congenital hemophilia A (without inhibitors) <input type="checkbox"/> D68.2 - Other congenital factor deficiency (FXIII) <input type="checkbox"/> D66 - Congenital hemophilia A (with inhibitors) <input type="checkbox"/> D68.311 - Acquired hemophilia <input type="checkbox"/> D67 - Congenital hemophilia B (without inhibitors) <input type="checkbox"/> D69.1 - Qualitative platelet defect (Glanzmann's Thrombasthenia) <input type="checkbox"/> D67 - Congenital hemophilia B (with inhibitors) Other diagnosis: <input type="checkbox"/> D68.2 - Other congenital factor deficiency (FVII) ICD-10 code and description: _____

MEDICAL ASSESSMENT	Patient Weight (kg): _____
	IV Access: <input type="checkbox"/> PIV/butterfly <input type="checkbox"/> Implanted Port <input type="checkbox"/> PICC <input type="checkbox"/> Central Line
	Additional Information: _____

PRESCRIPTION	Order Information: Complete prescription information below or submit a prescription with the strengths and assay limits. Quantity limits apply.				
	Product name:	Dose:	Directions:	Qty:	Refills:
	Highlights of prescribing information on following pages. For full prescribing information, see product specific package insert.				Do you intend to buy and bill? <input type="checkbox"/> Yes <input type="checkbox"/> No
	Preferred pharmacy:	Pharmacy Phone:		Pharmacy Fax:	
Pharmacy address:		City:	State:	Zip:	

PRESCRIBER AUTHORIZATION	Prescriber name:*		License #:*		
	Practice name and office contact:				Preferred method of contact: <input type="checkbox"/> Phone <input type="checkbox"/> Fax <input type="checkbox"/> Email
	DEA #:	Tax ID #:		NPI #:*	
	Phone:*	Fax:*	Email:*		
	Address:*		City:*	State:*	Zip:*
	Prescriber release: * By signing below, I hereby certify that: (a) I am a licensed practitioner, in good standing under applicable state law; (b) in my medical judgment, I have determined that the product(s) being prescribed which are detailed on the accompanying prescribing information (and listed only with those on label indications, appropriate treatment type, approved patient population, labeled dosing, and frequency) is to treat a diagnosis(es) consistent with indications dosing, and appropriate uses as described in the product's prescribing information; (c) the information I have provided on this enrollment form is, to the best of my knowledge, true, complete, and accurate in all respects; and (d) I have obtained the necessary authorization from the patient, or where appropriate the patient's parent, caregiver, and/or legal representative to use, disclose, share, and/or release the above-referenced information along with other protected health information (as defined in the Health Insurance Portability and Accountability Act of 1996 ("HIPAA")) for the sole purpose of providing patient assistance. Further, I appoint NovoCare [®] , on my behalf, to convey this prescription to the dispensing pharmacy. I will immediately notify Novo Nordisk Inc., its employees, or partners, including AssistRx, Inc. (collectively, "NovoCare [®] ") if the above-named patient, or where appropriate the patient's parent, caregiver, and/or legal representative, revokes their consent to share their PHI with NovoCare [®] . I give you permission to contact me with any questions related to NovoCare [®] .				

Prescriber signature (no signature stamps):*				Date:*
---	--	--	--	--------



RARE BLOOD DISORDERS | HIGHLIGHTS OF PRESCRIBING INFORMATION

Novoeight[®]: Hemophilia A (congenital FVIII deficiency)

Treatment type	Patient Population/Bleed type	Dose/Target factor level	Frequency
Prophylaxis	Adult (≥ 12 yrs)	20-50 IU/kg	3 times weekly
	Adult (≥ 12 yrs)	20-40 IU/kg	Every other day
	Pediatric	25-60 IU/kg	3 times weekly
	Pediatric	25-50 IU/kg	Every other day
On-demand	Minor bleed	20-40 IU/dL	Every 12-24 hours
	Moderate bleed	30-60 IU/dL	Every 12-24 hours
	Major bleed	60-100 IU/dL	Every 8-24 hours
Perioperative	Minor surgery	30-60 IU/dL	Every 24 hours
	Major surgery	80-100 IU/dL	Every 8-24 hours

Esperoct[®]: Hemophilia A (congenital FVIII deficiency)

Treatment type	Patient Population/Bleed type	Dose	Frequency
Prophylaxis	Adult (≥ 12 yrs)	50 IU/kg	Every 4 days*
	Pediatric (< 12 years)	65 IU/kg	2 times weekly*
On-demand	Adult: Minor bleed	40 IU/kg	1 dose should be sufficient
	Adult: Moderate bleed	40 IU/kg	An additional dose may be administered after 24 hours
	Adult: Major bleed	50 IU/kg	Additional dose(s) may be administered approximately every 24 hours
	Pediatric: any bleed	65 IU/kg	Minor: 1 dose should be sufficient Moderate: An additional dose may be administered after 24 hours Major: Additional dose(s) may be administered approximately every 24 hours
Perioperative	Adult: Minor or Major	50 IU/kg	Minor: Every 24 hours
			Major: Every 24 hours for the first week, then approximately every 48 hours until wound healing has occurred
	Pediatric: Minor or Major	65 IU/kg	Minor: Every 24 hours
			Major: Every 24 hours for the first week, then approximately every 48 hours until wound healing has occurred

*Frequency can be adjusted based on bleeding episodes

Rebinyn[®]: Hemophilia B (congenital FIX deficiency); All doses/frequencies are for both adult and pediatric populations

Treatment type	Patient Population/Bleed type	Dose	Frequency
Prophylaxis	N/A	40 IU/kg	Once weekly*
On-demand	Minor/Moderate bleed	40 IU/kg	1 dose should be sufficient, but additional doses of 40 IU/kg can be given
	Major bleed	80 IU/kg	Additional doses of 40 IU/kg can be given
Perioperative	Minor surgery	40 IU/kg	1 pre-op dose should be sufficient, additional doses of 40 IU/kg can be given
	Major surgery	80 IU/kg	Pre-op dose; additional doses of 40 IU/kg can be given every 1-3 days within 1st week

*Frequency can be adjusted based on bleeding episodes and physical activity

Patient Enrollment Form | Rare Blood Disorders

Phone: 1-844-668-6732
 Fax: 1-866-488-6576

Monday - Friday
 8:00 AM to 8:00 PM ET



NovoCare[®]
 Savings | Coverage | Support

Tretten[®]: Congenital FXIII A-subunit deficiency; adult and pediatric populations

Treatment type	Patient Population/Bleed type	Dose	Frequency
Prophylaxis	N/A	35 IU/kg	Monthly*

*Consider dose adjustment if adequate coverage is not achieved

NovoSeven[®] RT

Indication	Treatment type	Patient Population/Bleed type	Dose	Frequency
Hemophilia A with inhibitors or hemophilia B with inhibitors	On-demand	Adult/pediatric: all other bleeds	90 mcg/kg	Every 2 hours until hemostasis is achieved, or until the treatment has been judged to be inadequate
		Adult/pediatric: severe bleeds	90 mcg/kg	Every 2 hours until hemostasis and then post hemostatic every 3-6 hours
	Perioperative	Adult/pediatric: minor surgery	90 mcg/kg	Immediately before surgery, every 2 hours during surgery, every 2 hours after surgery for 48 hours and then every 2-6 hours until healing occurs
		Adult/pediatric: major surgery	90 mcg/kg	Immediately before surgery, every 2 hours during surgery, every 2 hours after surgery for 5 days and then every 4 hours or by continuous infusion (50 mcg/kg/hr) until healing occurs
Congenital FVII deficiency	On-demand	Adult/pediatric	15-30 mcg/kg	Every 4-6 hours until hemostasis achieved
	Perioperative	Adult/pediatric	15-30 mcg/kg	Immediately before surgery, every 4-6 hours during surgery and until healing occurs
Glanzmann's thrombasthenia (with refractoriness to platelet transfusions)	On-demand	Adult/pediatric	90 mcg/kg	Every 2-6 hours until hemostasis achieved
	Perioperative	Adult/pediatric	90 mcg/kg	Immediately before surgery, every 2 hours during surgery and every 2-6 hours post surgery
Acquired hemophilia	On-demand	Adult	70-90 mcg/kg	Every 2-3 hours until hemostasis achieved
	Perioperative	Adult	70-90 mcg/kg	Immediately before surgery, every 2-3 hours during surgery and until hemostasis achieved